

## AQUA 27: APPROPRIATE TESTING FOR VASECTOMY PATIENTS

## STEWARD

AMERICAN UROLOGICAL ASSOCIATION

## DESCRIPTION

Percentage of patients where a Post Vasectomy Semen Analysis (PVSA) was ordered and confirmed sterility within 6 months of undergoing a vasectomy

## TYPE

Process

## DATA SOURCE

Electronic Health Records: AQUA Registry Data

## NUMERATOR STATEMENT

Patients with an order for a PVSA 6 months after a vasectomy and confirmed sterility (fresh uncentrifuged semen sample performed)

## DENOMINATOR STATEMENT

All patients undergoing a vasectomy

## DENOMINATOR DETAILS

List Denominator Codes

## DENOMINATOR EXCLUSIONS/EXCEPTIONS

Patients with known renal insufficiency (Cr >1.5 or documented in past medical history) or with documented flank pain or hematuria

## RATIONALE

PVSA testing should be performed post-vasectomy. Approximately 15% of patients with azoospermia will have a genetic etiology and thus these patients should have a y-chromosome microdeletion assessment as well as a karyotype.

## RISK ADJUSTMENT

No

## INVERSE MEASURE

Yes

## ALGORITHM

## COPYRIGHT/DISCLAIMER

AMERICAN UROLOGICAL ASSOCIATION