

AQUA 28: HYPOGONADISM: TESTOSTERONE AND HEMATOCRIT WITHIN 6 MONTHS OF STARTING TESTOSTERONE REPLACEMENT

STEWARD

AMERICAN UROLOGICAL ASSOCIATION

DESCRIPTION

Percentage of patients with hypogonadism, starting testosterone medication (any formulation of testosterone) or received testosterone injection in clinic have testosterone level ordered within 6 months of first testosterone Rx or injection

TYPE

Process

DATA SOURCE

Electronic Health Records: AQUA Registry Data

NUMERATOR STATEMENT

Number of patients with a diagnosis of hypogonadism receiving testosterone replacement (medication or injection) with a testosterone level test and hematocrit ordered/reported within 6 months of first testosterone Rx or infection

DENOMINATOR STATEMENT

Number of patients with a diagnosis of hypogonadism receiving testosterone replacement (medication or injection)

DENOMINATOR DETAILS

List Denominator Codes

DENOMINATOR EXCLUSIONS/EXCEPTIONS

RATIONALE

Measure reflects the most effective prevention and treatment processes

RISK ADJUSTMENT

No

INVERSE MEASURE

No

ALGORITHM

COPYRIGHT/DISCLAIMER

AMERICAN UROLOGICAL ASSOCIATION