

## Non-QPP Measures

#	Measure ID	Measure Title	Definition	Historical Benchmark <sup>†</sup>	Type	High Priority Measure	Included in Supported MVPs
1	<b>AQUA8®</b>	Hospital Admissions or Infectious Complications Within 30 days of Prostate Biopsy	Percentage of patients with urinary retention, infection, inpatient consultation, or hospital admission for infection or sepsis within 30 days of undergoing a prostate biopsy	Yes	Outcome	Yes	Optimal Care for Patients with Urologic Conditions
2	<b>AQUA14®</b>	Stones: Repeat Shock Wave Lithotripsy (SWL) Within 6 Months of Initial Treatment	Percentage of patients who underwent repeat Shock Wave Lithotripsy within 6 months of initial procedure	Yes	Outcome	Yes	Optimal Care for Patients with Urologic Conditions
3	<b>AQUA15®</b>	Stones: Urinalysis or Urine Culture Performed Before Surgical Stone Procedures	Percentage of patients with a documented urinalysis or urine culture within 14 days prior to surgical stone procedures	Yes	Process	Yes	Optimal Care for Patients with Urologic Conditions
4	<b>AQUA16®</b>	Non-Muscle Invasive Bladder Cancer: Repeat Transurethral Resection of Bladder Tumor (TURBT) for T1 disease	Percentage of patients with T1 disease who had a second transurethral resection of bladder tumor (TURBT) within 6 weeks of the initial TURBT	No	Process	No	Optimal Care for Patients with Urologic Conditions
5	<b>AQUA35®</b>	Non-Muscle Invasive Bladder Cancer: Initial Management/Surveillance for Non-Muscle Invasive Bladder Cancer	Percentage of patients with appropriate initial management/surveillance after initial diagnosis of non-muscle invasive bladder cancer	No (new in 2024: possible 5-point floor)	Process	No	N/A

<sup>†</sup>Based on 2/5/2025 benchmark file from CMS for PY2025

\*Electronic Clinical Quality Measure

\*©2011 The Regents of the University of Michigan. Original MUSIC measure developed by the University of Michigan as part of the Michigan Urological Surgery Improvement Collaborative. All Rights Reserved.



## Non-QPP Measures

#	Measure ID	Measure Title	Definition	Historical Benchmark <sup>†</sup>	Type	High Priority Measure	Included in Supported MVPs
6	<b>AQUA36<sup>®</sup></b>	Prostate Cancer: Confirmation Biopsy in Newly Diagnosed Patients on Active Surveillance	Percentage of newly diagnosed low risk patients on active surveillance who receive confirmation biopsy within 24 months of diagnosis	No (new in 2024: possible 5-point floor)	Process	No	N/A
7	<b>MUSIC4<sup>‡</sup></b>	Prostate Cancer: Active Surveillance/Watchful Waiting for Newly Diagnosed Low Risk Prostate Cancer Patients	Proportion of patients newly diagnosed with low-risk prostate cancer managed with active surveillance or watchful waiting	No	Process	Yes	Optimal Care for Patients with Urologic Conditions

<sup>†</sup>Based on 2/5/2025 benchmark file from CMS for PY2025

<sup>\*</sup>Electronic Clinical Quality Measure

<sup>‡</sup>©2011 The Regents of the University of Michigan. Original MUSIC measure developed by the University of Michigan as part of the Michigan Urological Surgery Improvement Collaborative. All Rights Reserved.

## QPP Measures

#	QPP #	Measure Title	Definition	Historical Benchmark <sup>†</sup>	Type	High Priority Measure	Included in Supported MVPs
8	001	Diabetes: Glycemic Status Assessment Greater Than 9%	Percentage of patients 18-75 years of age with diabetes who had a glycemic status assessment (hemoglobin A1c [HbA1c] or glucose management indicator [GMI]) > 9.0% during the measurement period.	Yes	Intermediate Outcome	Yes	N/A
9	eCQM* CMS122v13			Yes			
10	047	Advance Care Plan	Percentage of patients aged 65 years and older who have an advance care plan or surrogate decision maker documented in the medical record or documentation in the medical record that an advance care plan was discussed but the patient did not wish or was not able to name a surrogate decision maker or provide an advance care plan.	Yes	Process	Yes	Advancing Cancer Care
11	048	Urinary Incontinence: Assessment of Presence or Absence of Urinary Incontinence in Women Aged 65 Years and Older	Percentage of female patients aged 65 years and older who were assessed for the presence or absence of urinary incontinence within 12 months.	Yes	Process	No	Focusing on Women's Health
12	050	Urinary Incontinence: Plan of Care for Urinary Incontinence in Women Aged 65 Years and Older	Percentage of female patients aged 65 years and older with a diagnosis of urinary incontinence with a documented plan of care for urinary incontinence at least once within 12 months.	Yes	Process	Yes	Optimal Care for Patients with Urologic Conditions

<sup>†</sup>Based on 2/5/2025 benchmark file from CMS for PY2025

\*Electronic Clinical Quality Measure

<sup>†</sup>©2011 The Regents of the University of Michigan. Original MUSIC measure developed by the University of Michigan as part of the Michigan Urological Surgery Improvement Collaborative. All Rights Reserved.

## QPP Measures

#	QPP #	Measure Title	Definition	Historical Benchmark <sup>†</sup>	Type	High Priority Measure	Included in Supported MVPs
13	102	Prostate Cancer: Avoidance of Overuse of Bone Scan for Staging Low Risk Prostate Cancer Patients	Percentage of patients, regardless of age, with a diagnosis of prostate cancer at low (or very low) risk of recurrence receiving interstitial prostate brachytherapy, OR external beam radiotherapy to the prostate, OR radical prostatectomy who did not have a bone scan performed at any time since diagnosis of prostate cancer.	No	Process	Yes	Advancing Cancer Care
14	eCQM* CMS129v14			No			
15	130	Documentation of Current Medications in the Medical Record	Percentage of visits for which the eligible clinician attests to documenting a list of current medications using all immediate resources available on the date of the encounter.	Yes	Process	Yes	N/A
16	eCQM* CMS68v14			Yes			
17	134	Preventive Care and Screening: Screening for Depression and Follow-Up Plan	Percentage of patients aged 12 years and older screened for depression on the date of the encounter or up to 14 days prior to the date of the encounter using an age-appropriate standardized depression screening tool AND if positive, a follow-up plan is documented on the date of or up to two days after the date of the qualifying encounter.	Yes	Process	No	Advancing Cancer Care; Focusing on Women's Health
18	eCQM* CMS2v14			Yes			

<sup>†</sup>Based on 2/5/2025 benchmark file from CMS for PY2025

\*Electronic Clinical Quality Measure

<sup>‡</sup>©2011 The Regents of the University of Michigan. Original MUSIC measure developed by the University of Michigan as part of the Michigan Urological Surgery Improvement Collaborative. All Rights Reserved.

## QPP Measures

#	QPP #	Measure Title	Definition	Historical Benchmark <sup>†</sup>	Type	High Priority Measure	Included in Supported MVPs
19	143	Oncology: Medical and Radiation - Pain Intensity Quantified	Percentage of patient visits, regardless of patient age, with a diagnosis of cancer currently receiving chemotherapy or radiation therapy in which pain intensity is quantified.	Yes	Process	Yes	Advancing Cancer Care
20	eCQM* CMS157v13			Yes			
21	144	Oncology: Medical and Radiation - Plan of Care for Pain	Percentage of visits for patients, regardless of age, with a diagnosis of cancer currently receiving chemotherapy or radiation therapy who report having pain with a documented plan of care to address pain.	Yes	Process	Yes	Advancing Cancer Care

<sup>†</sup>Based on 2/5/2025 benchmark file from CMS for PY2025

\*Electronic Clinical Quality Measure

<sup>‡</sup>©2011 The Regents of the University of Michigan. Original MUSIC measure developed by the University of Michigan as part of the Michigan Urological Surgery Improvement Collaborative. All Rights Reserved.

## QPP Measures

#	QPP #	Measure Title	Definition	Historical Benchmark <sup>†</sup>	Type	High Priority Measure	Included in Supported MVPs
22	226	Preventive Care and Screening: Tobacco Use: Screening and Cessation Intervention	Percentage of patients aged 12 years and older who were screened for tobacco use one or more times within the measurement period and who received tobacco cessation intervention during the measurement period or in the six months prior to the measurement period if identified as a tobacco user.	Yes	Process	No	Focusing on Women's Health
23	eCQM* CMS138v13			Yes			
24	236	Controlling High Blood Pressure	Percentage of patients 18-85 years of age who had a diagnosis of essential hypertension starting before and continuing into, or starting during the first six months of the measurement period, and whose most recent blood pressure was adequately controlled (<140/90mmHg) during the measurement period.	Yes	Intermediate Outcome	Yes	N/A
25	eCQM* CMS165v13			Yes			

<sup>†</sup>Based on 2/5/2025 benchmark file from CMS for PY2025

\*Electronic Clinical Quality Measure

<sup>†</sup>©2011 The Regents of the University of Michigan. Original MUSIC measure developed by the University of Michigan as part of the Michigan Urological Surgery Improvement Collaborative. All Rights Reserved.

## QPP Measures

#	QPP #	Measure Title	Definition	Historical Benchmark <sup>†</sup>	Type	High Priority Measure	Included in Supported MVPs
26	238	Use of High-Risk Medications in Older Adults	Percentage of patients 65 years of age and older who were ordered at least two high-risk medications from the same drug class.	Yes	Process	Yes	N/A
27	eCQM* CMS156v13			Yes			
28	317	Preventive Care and Screening: Screening for High Blood Pressure and Follow-Up Documented	Percentage of patient visits for patients aged 18 years and older seen during the performance period who were screened for high blood pressure AND a recommended follow-up plan is documented, as indicated, if blood pressure is elevated or hypertensive.	Yes	Process	No	N/A
29	eCQM* CMS22v13			Yes			
30	318 eCQM* CMS139v13 version only	Falls: Screening for Future Fall Risk	Percentage of patients 65 years of age and older who were screened for future fall risk during the measurement period.	Yes	Process	Yes	Optimal Care for Patients with Urologic Conditions
31	357	Surgical Site Infection (SSI)	Percentage of patients aged 18 years and older who had a surgical site infection (SSI).	Yes	Outcome	Yes	N/A

<sup>†</sup>Based on 2/5/2025 benchmark file from CMS for PY2025

\*Electronic Clinical Quality Measure

<sup>†</sup>©2011 The Regents of the University of Michigan. Original MUSIC measure developed by the University of Michigan as part of the Michigan Urological Surgery Improvement Collaborative. All Rights Reserved.

## QPP Measures

#	QPP #	Measure Title	Definition	Historical Benchmark <sup>†</sup>	Type	High Priority Measure	Included in Supported MVPs
32	358	Patient-Centered Surgical Risk Assessment and Communication	Percentage of patients who underwent a non-emergency surgery who had their personalized risks of postoperative complications assessed by their surgical team prior to surgery using a clinical data-based, patient-specific risk calculator and who received personal discussion of those risks with the surgeon.	Yes	Process	Yes	Optimal Care for Patients with Urologic Conditions
33	370	Depression Remission at Twelve Months	The percentage of adolescent patients 12 to 17 years of age and adult patients 18 years of age or older with major depression or dysthymia who reached remission 12 months (+/- 60 days) after an index event date.	No	Outcome	Yes	N/A
34	eCQM* CMS159v13			Yes			
35	374	Closing the Referral Loop: Receipt of Specialist Report	Percentage of patients with referrals, regardless of age, for which the referring clinician receives a report from the clinician to whom the patient was referred.	Yes	Process	Yes	N/A
36	eCQM* CMS50v13			Yes			

<sup>†</sup>Based on 2/5/2025 benchmark file from CMS for PY2025

\*Electronic Clinical Quality Measure

\*©2011 The Regents of the University of Michigan. Original MUSIC measure developed by the University of Michigan as part of the Michigan Urological Surgery Improvement Collaborative. All Rights Reserved.



## QPP Measures

#	QPP #	Measure Title	Definition	Historical Benchmark <sup>†</sup>	Type	High Priority Measure	Included in Supported MVPs
37	422	Performing Cystoscopy at the Time of Hysterectomy for Pelvic Organ Prolapse to Detect Lower Urinary Tract Injury	Percentage of patients who undergo cystoscopy to evaluate for lower urinary tract injury at the time of hysterectomy for pelvic organ prolapse.	No	Process	Yes	Focusing on Women's Health
38	431	Preventive Care and Screening: Unhealthy Alcohol Use: Screening & Brief Counseling	Percentage of patients aged 18 years and older who were screened for unhealthy alcohol use using a systematic screening method at least once within the last 12 months AND who received brief counseling if identified as an unhealthy alcohol user.	Yes	Process	No	Focusing on Women's Health
39	432	Proportion of Patients Sustaining a Bladder or Bowel Injury at the time of any Pelvic Organ Prolapse Repair	Percentage of patients undergoing surgical repair of pelvic organ prolapse that is complicated by a bladder or bowel injury at the time of index surgery that is recognized intraoperatively or within 30 days after surgery.	No	Outcome	Yes	Focusing on Women's Health
40	453	Percentage of Patients who Died from Cancer Receiving Systemic Cancer-Directed Therapy in the Last 14 Days of Life (lower score – better)	Percentage of patients who died from cancer receiving systemic cancer-directed therapy in the last 14 days of life.	Yes	Process	Yes	Advancing Cancer Care

<sup>†</sup>Based on 2/5/2025 benchmark file from CMS for PY2025

\*Electronic Clinical Quality Measure

\*©2011 The Regents of the University of Michigan. Original MUSIC measure developed by the University of Michigan as part of the Michigan Urological Surgery Improvement Collaborative. All Rights Reserved.

## QPP Measures

#	QPP #	Measure Title	Definition	Historical Benchmark <sup>†</sup>	Type	High Priority Measure	Included in Supported MVPs
41	457	Percentage of Patients Who Died from Cancer Admitted to Hospice for Less than 3 days (lower score – better)	Percentage of patients who died from cancer, and admitted to hospice and spent less than 3 days there.	Yes	Process	Yes	Advancing Cancer Care
42	462 <i>eCQM*</i> <i>CMS645v8</i> <i>version only</i>	Bone Density Evaluation for Patients with Prostate Cancer and Receiving Androgen Deprivation Therapy	Percentage of patients determined as having prostate cancer who are currently starting or undergoing androgen deprivation therapy (ADT), for an anticipated period of 12 months or greater and who receive an initial bone density evaluation. The bone density evaluation must be prior to the start of ADT or within 3 months of the start of ADT.	No	Process	No	Advancing Cancer Care; Optimal Care for Patients with Urologic Conditions
43	476 <i>eCQM*</i> <i>CMS771v6</i> <i>version only</i>	Urinary Symptom Score Change 6-12 Months After Diagnosis of Benign Prostatic Hyperplasia	Percentage of patients with an office visit within the measurement period and with a new diagnosis of clinically significant Benign Prostatic Hyperplasia who have International Prostate Symptom Score (IPSS) or American Urological Association (AUA) Symptom Index (SI) documented at time of diagnosis and again 6-12 months later with an improvement of 3 points.	No	Patient-Reported Outcome-Based Performance Measure	Yes	Optimal Care for Patients with Urologic Conditions

<sup>†</sup>Based on 2/5/2025 benchmark file from CMS for PY2025

\*Electronic Clinical Quality Measure

<sup>‡</sup>©2011 The Regents of the University of Michigan. Original MUSIC measure developed by the University of Michigan as part of the Michigan Urological Surgery Improvement Collaborative. All Rights Reserved.

## QPP Measures

#	QPP #	Measure Title	Definition	Historical Benchmark <sup>†</sup>	Type	High Priority Measure	Included in Supported MVPs
44	<b>481</b> <b>eCQM*</b> <b>CMS646v5</b> <b>version only</b>	Intravesical Bacillus-Calmette-Guerin for Non-Muscle Invasive Bladder Cancer	Percentage of patients initially diagnosed with non-muscle invasive bladder cancer and who received intravesical Bacillus-Calmette-Guerin (BCG) within 6 months of bladder cancer staging.	No	Process	Yes	Optimal Care for Patients with Urologic Conditions
45	<b>483</b>	Person-Centered Primary Care Measure Patient Reported Outcome Performance Measure (PCPCM PRO-PM)	The Person-Centered Primary Care Measure Patient Reported Outcome Performance Measure (PCPCM PRO-PM) uses the PCPCM Patient Reported Outcome Measure (PROM) a comprehensive and parsimonious set of 11 patient-reported items - to assess the broad scope of primary care. Unlike other primary care measures, the PCPCM PRO-PM measures the high value aspects of primary care based on a patient's relationship with the clinician or practice.	No	Patient-Reported Outcome-Based Performance Measure	Yes	N/A

<sup>†</sup>Based on 2/5/2025 benchmark file from CMS for PY2025

\*Electronic Clinical Quality Measure

<sup>‡</sup>©2011 The Regents of the University of Michigan. Original MUSIC measure developed by the University of Michigan as part of the Michigan Urological Surgery Improvement Collaborative. All Rights Reserved.

## QPP Measures

#	QPP #	Measure Title	Definition	Historical Benchmark <sup>†</sup>	Type	High Priority Measure	Included in Supported MVPs
46	487	Screening for Social Drivers of Health	Percent of patients 18 years and older screened for food insecurity, housing instability, transportation needs, utility difficulties, and interpersonal safety.	Yes	Process	Yes	Advancing Cancer Care; Focusing on Women's Health; Optimal Care for Patients with Urologic Conditions
47	488	Kidney Health Evaluation	Percentage of patients aged 18-85 years with a diagnosis of diabetes who received a kidney health evaluation defined by an Estimated Glomerular Filtration Rate (eGFR) AND Urine Albumin-Creatinine Ratio (uACR) within the measurement period.	Yes	Process	No	N/A
48	eCQM* CMS951v3			No			

<sup>†</sup>Based on 2/5/2025 benchmark file from CMS for PY2025

\*Electronic Clinical Quality Measure

<sup>‡</sup>©2011 The Regents of the University of Michigan. Original MUSIC measure developed by the University of Michigan as part of the Michigan Urological Surgery Improvement Collaborative. All Rights Reserved.

## QPP Measures

#	QPP #	Measure Title	Definition	Historical Benchmark <sup>†</sup>	Type	High Priority Measure	Included in Supported MVPs
49	<b>490</b>	Appropriate Intervention of Immune-Related Diarrhea and/or Colitis in Patients Treated with Immune Checkpoint Inhibitors	Percentage of patients, aged 18 years and older, with a diagnosis of cancer, on immune checkpoint inhibitor therapy, and grade 2 or above diarrhea and/or grade 2 or above colitis, who have immune checkpoint inhibitor therapy held and corticosteroids or immunosuppressants prescribed or administered.	No	Process	No	Advancing Cancer Care
50	<b>493</b>	Adult Immunization Status	Percentage of patients 19 years of age and older who are up-to-date on recommended routine vaccines for influenza; tetanus and diphtheria (Td) or tetanus, diphtheria and acellular pertussis (Tdap); zoster; and pneumococcal.	Yes	Process	No	Focusing on Women's Health

<sup>†</sup>Based on 2/5/2025 benchmark file from CMS for PY2025

\*Electronic Clinical Quality Measure

<sup>‡</sup>©2011 The Regents of the University of Michigan. Original MUSIC measure developed by the University of Michigan as part of the Michigan Urological Surgery Improvement Collaborative. All Rights Reserved.

## QPP Measures

#	QPP #	Measure Title	Definition	Historical Benchmark <sup>†</sup>	Type	High Priority Measure	Included in Supported MVPs
51	497	Preventive Care and Wellness (Composite)	Percentage of patients who received age- and sex-appropriate preventive screenings and wellness services. This measure is a composite of seven component measures that are based on recommendations for preventive care by the U.S. Preventive Services Task Force (USPSTF), Advisory Committee on Immunization Practices (ACIP), American Association of Clinical Endocrinology (AACE), and American College of Endocrinology (ACE).	No (new in 2024: possible 5-point floor)	Process	No	N/A
52	498	Connection to Community Service Provider	Percent of patients 18 years or older who screen positive for one or more of the following health-related social needs (HRSNs): food insecurity, housing instability, transportation needs, utility help needs, or interpersonal safety; and had contact with a Community Service Provider (CSP) for at least one of their HRSNs within 60 days after screening.	No (new in 2024: possible 5-point floor)	Process	Yes	N/A

<sup>†</sup>Based on 2/5/2025 benchmark file from CMS for PY2025

\*Electronic Clinical Quality Measure

<sup>‡</sup>©2011 The Regents of the University of Michigan. Original MUSIC measure developed by the University of Michigan as part of the Michigan Urological Surgery Improvement Collaborative. All Rights Reserved.

## QPP Measures

#	QPP #	Measure Title	Definition	Historical Benchmark <sup>†</sup>	Type	High Priority Measure	Included in Supported MVPs
53	503	Gains in Patient Activation Measure (PAM) Scores at 12 Months	The Patient Activation Measure® (PAM®) is a 10- or 13-item questionnaire that assesses an individual's knowledge, skills, and confidence for managing their health and health care. The measure assesses individuals on a 0-100 scale that converts to one of four levels of activation, from low (1) to high (4). The PAM® performance measure (PAM®-PM) is the change in score on the PAM® from baseline to follow-up measurement.	No (new in 2024: possible 5-point floor)	Patient-Reported Outcome-Based Performance Measure	Yes	Advancing Cancer Care; Optimal Care for Patients with Urologic Conditions
54	508	Adult COVID-19 Vaccination Status	Percentage of patients aged 18 years and older seen for a visit during the performance period that are up-to-date on their COVID-19 vaccinations as defined by Centers for Disease Control and Prevention (CDC) recommendations on current vaccination.	No (new in 2025: possible 7-point floor)	Process	No	N/A

<sup>†</sup>Based on 2/5/2025 benchmark file from CMS for PY2025

\*Electronic Clinical Quality Measure

\*©2011 The Regents of the University of Michigan. Original MUSIC measure developed by the University of Michigan as part of the Michigan Urological Surgery Improvement Collaborative. All Rights Reserved.