

**Non-QPP Measures**

#	Measure ID	Measure Title	Definition	Historical Benchmark?	Type	Domain
1	AQUA8 <sup>^</sup>	Hospital admissions/complications within 30 days of TRUS Biopsy	Percentage of patients who had TRUS biopsy performed who had ≥24h after the biopsy): infection, hematuria, new antibiotic Rx after biopsy, or inpatient consultation within 30 days	Yes	Outcome	Patient Safety
2	AQUA14 <sup>^</sup> (inverse)	Stones: Inappropriate Repeat Shock Wave Lithotripsy (SWL) Within 6 Months of Initial Treatment	Percentage of patients who underwent inappropriate repeat Shock Wave Lithotripsy within 6 months of initial procedure	No	Outcome	Efficiency and Cost Reduction
3	AQUA15 <sup>^</sup>	Stones: Urinalysis Performed Before Surgical Stone Procedures	Percentage of patients with a documented urinalysis 30 days before surgical stone procedures	Yes	Process	Patient Safety
4	AQUA16	Non-Muscle Invasive Bladder Cancer: Repeat Transurethral Resection of Bladder Tumor (TURBT) for T1 disease	Percentage of patients with T1 disease, that had a second TURBT within 6 weeks of the initial TURBT	No	Process	Efficiency and Cost Reduction
5	AQUA18	Non-Muscle Invasive Bladder Cancer: Early Surveillance Cystoscopy for Non-Muscle Invasive Bladder Cancer	Percentage of patients who receive surveillance cystoscopy within 4 months of TURBT for bladder cancer	Yes	Process	Effective Clinical Care
6	AQUA26 <sup>^</sup> (inverse)	Benign Prostate Hyperplasia (BPH): Inappropriate Lab & Imaging Services for Patients with BPH	Percentage of patients with new diagnosis of BPH who had a creatinine lab order placed or had a CT abdomen, MRI abdomen, ultrasound abdomen ordered or performed.	Yes	Process	Efficiency and Cost Reduction

**Non-QPP Measures**

#	Measure ID	Measure Title	Definition	Historical Benchmark?	Type	Domain
7	<b>MUSIC4</b>	Prostate Cancer: Active surveillance/watchful waiting for low-risk prostate cancer	Percentage of patients with low-risk prostate cancer receiving active surveillance or watchful waiting	No	Process	Effective Clinical Care
8	<b>MUSIC10</b>	Prostate Cancer: Confirmation Testing in low risk AS eligible patients	Percentage of low risk patients that are eligible for active surveillance who receive confirmation testing within 6 months of diagnosis	No	Process	Efficiency and Cost Reduction
9	<b>MUSIC11</b>	Prostate Cancer: Follow-Up Testing for patients on active surveillance for at least 30 months	Percentage of patients on active surveillance that have $\geq 2$ tumor burden reassessments and 3 PSA tests in first 30 months since diagnosis	Yes	Process	Effective Clinical Care

**QPP Measures**

#	QPP #	Measure Title	Definition	Historical Benchmark?	Type	Domain
10	1 <sup>^</sup>	Diabetes: Hemoglobin A1c (HbA1c) Poor Control (>9%)	Percentage of patient 18-75 years of age with diabetes who had hemoglobin A1c>9.0% during the measurement period	Yes	Intermediate Outcome	Effective Clinical Care
11	eCQM* CMS122v8			Yes		
12	23 <sup>^</sup>	Perioperative Care: Venous Thromboembolism (VTE) Prophylaxis (When indicated in ALL patients)	Percentage of surgical patients aged 18 years and older undergoing procedures for which VTE prophylaxis is indicated in all patients, who had an order for Low Molecular Weight Heparin (LMWH), Low-Dose Unfractionated Heparin (LDUH), adjusted-dose warfarin, fondaparinux or mechanical prophylaxis to be given within 24 hours prior to incision time or within 24 hours after	Yes	Process	Patient Safety
13	47 <sup>^</sup>	Advance Care Plan	Percentage of patients aged 65 years and older who have an advance care plan or surrogate decision maker documented in the medical record or documentation in the medical record that an advance care plan was discussed but the patient did not wish or was not able to name a surrogate decision maker or provide an advance care plan	Yes	Process	Communication &Care Coordination

**QPP Measures**

#	QPP #	Measure Title	Definition	Historical Benchmark?	Type	Domain
14	48	Urinary Incontinence: Assessment of Presence or Absence of Urinary Incontinence in Women Aged 65 Years and Older	Percentage of female patients aged 65 years and older who were assessed for the presence or absence of urinary incontinence within 12 months	Yes	Process	Effective Clinical Care
15	50^	Urinary Incontinence: Plan of Care for Urinary Incontinence in Women Aged 65 Years and Older	Percentage of female patients aged 65 years and older with a diagnosis of urinary incontinence with a documented plan of care for urinary incontinence at least once within 12 months	Yes	Process	Person & Caregiver-Centered Experience & Outcomes
16	102^	Prostate Cancer: Avoidance of Overuse of Bone Scan for Staging Low Risk Prostate Cancer Patients	Percentage of patients, regardless of age, with a diagnosis of prostate cancer at low risk of recurrence receiving interstitial prostate brachytherapy, OR external beam radiotherapy to the prostate, OR radical prostatectomy, OR cryotherapy who did not have a bone scan performed at any time since diagnosis of prostate cancer	Yes	Process	Efficiency and Cost Reduction
17	eCQM* CMS129v9			Yes		
18	104	Prostate Cancer: Combination Androgen Deprivation Therapy for High Risk or Very High Risk Prostate Cancer	Percentage of patients, regardless of age, with a diagnosis of prostate cancer at high risk of recurrence receiving external beam radiotherapy to the prostate who were prescribed adjuvant hormonal therapy (GnRH agonist or antagonist)	Yes	Process	Effective Clinical Care
19	110	Preventative Care and Screening: Influenza Immunization	Percentage of patients aged 6 months and older seen for a visit between October 1 and March 31 who received an influenza immunization OR who reported previous receipt of an influenza immunization	Yes	Process	Community/ Population Health
20	eCQM* CMS147v9			Yes		

**QPP Measures**

#	QPP #	Measure	Definition	Historical Benchmark	Type	Domain
21	111	Pneumococcal Vaccination Status for Older Adults	Percentage of patients 65 years of age and older who have ever received a pneumococcal vaccine	Yes	Process	Community/ Population Health
22	eCQM* CMS127v8			Yes		
23	112	Breast Cancer Screening	Percentage of women 50-74 years of age who had a mammogram to screen for breast cancer	Yes	Process	Effective Clinical Care
24	eCQM* CMS125v8			Yes		
25	113	Colorectal Cancer Screening	Percentage of patients 50 through 75 years of age who had appropriate screening for colorectal cancer	Yes	Process	Effective Clinical Care
26	eCQM* CMS130v8			Yes		
27	117	Diabetes: Eye Exam	Percentage of patients 18 - 75 years of age with diabetes who had a retinal or dilated eye exam by an eyecare professional during the measurement period or a negative retinal or dilated eye exam (no evidence of retinopathy) in the 12 months prior to the measurement period	Yes	Process	Effective Clinical Care
28	eCQM* CMS131v8			Yes		
29	119	Diabetes: Medical Attention for Nephropathy	The percentage of patients 18-75 years of age with diabetes who had a nephropathy screening test or evidence of nephropathy during the measurement period	Yes	Process	Effective Clinical Care
30	eCQM* CMS134v8			Yes		
31	128	Preventive Care and Screening: Body Mass Index (BMI) Screening and Follow-Up Plan	Percentage of patients aged 18 years and older with a documented BMI during the current encounter or during the previous six months AND when the BMI is outside of normal parameters, a follow-up plan is documented during the encounter or during the previous six months of the encounter	Yes	Process	Community/ Population Health
32	eCQM* CMS69v8			Yes		

**QPP Measures**

#	QPP #	Measure Title	Definition	Historical Benchmark?	Type	Domain
33	130^	Documentation of Current Medications in the Medical Record	Percentage of specified visits for patients aged 18 years and older for which the eligible professional attests to documenting a list of current medications to the best of his/her knowledge and ability. This list must include ALL prescriptions, over-the- counters, herbals, and vitamin/mineral/dietary (nutritional) supplements AND must contain the medications' name, dosage, frequency and route of administration	Yes	Process	Patient Safety
34	eCQM* CMS68v9			Yes		
35	134	Preventive Care and Screening: Screening for Depression and Follow-Up Plan	Percentage of patients aged 12 years and older screened for depression on the date of the encounter using an age appropriate standardized depression screening tool AND if positive, a follow-up plan is documented on the date of the positive screen	Yes	Process	Community/ Population Health
36	eCQM* CMS2v9			Yes		
37	226	Preventive Care and Screening: Tobacco Use: Screening and Cessation Intervention	Percentage of patients aged 18 years and older who were screened for tobacco use one or more times within 24 months AND who received cessation counseling intervention if identified as a tobacco user	Yes	Process	Community/ Population Health
38	eCQM* CMS138v8			Yes		
39	236^	Controlling High Blood Pressure	Percentage of patients 18 through 85 years of age who had a diagnosis of hypertension and whose blood pressure was adequately controlled (< 140/90 mmHg) during the measurement period	Yes	Intermediate Outcome	Effective Clinical Care
40	eCQM* CMS165v8			Yes		

### QPP Measures

#	QPP #	Measure Title	Definition	Historical Benchmark?	Type	Domain
41	265 <sup>^</sup>	Biopsy Follow-Up	Percentage of new patients whose biopsy results have been reviewed and communicated to the primary care/referring physician and patient by the performing physician	Yes	Process	Communication & Care Coordination
42	317	Preventive Care and Screening: Screening for High Blood Pressure Follow-up Documented	Percentage of patients aged 18 years and older seen during the reporting period who were screened for high blood pressure AND a recommended follow-up plan is documented based on the current blood pressure (BP) reading as indicated	Yes	Process	Community/ Population Health
43	eCQM* CMS22v8			Yes		
44	318 <sup>^</sup> eCQM* CMS139v8 version only	Falls: Screening for Future Fall Risk	Percentage of patients 65 years of age and older who were screened for future fall risk during the measurement period	Yes	Process	Community/ Population Health
45	357 <sup>^</sup>	Surgical Site Infection	Percentage of patients aged 18 years and older who had a surgical site infection (SSI)	Yes	Outcome	Effective Clinical Care
46	358 <sup>^</sup>	Patient-Centered Surgical Risk Assessment and Communication	Percentage of patients who underwent a non-emergency surgery who had their personalized risks of postoperative complications assessed by their surgical team prior to surgery using a clinical data-based, patient-specific risk calculator	Yes	Process	Person & Caregiver-Centered Experience & Outcomes
47	370 <sup>^</sup>	Depression Remission at Twelve Months	The percentage of patients 18 years of age or older with major depression or dysthymia who reached remission 12 months (+/- 30 days) after an index visit	Yes	Outcome	Effective Clinical Care

**QPP Measures**

#	QPP #	Measure Title	Definition	Historical Benchmark?	Type	Domain
48	eCQM* CMS159v8	Depression Remission at Twelve Months	The percentage of patients 18 years of age or older with major depression or dysthymia who reached remission 12 months (+/- 30 days) after an index visit	Yes	Outcome	Effective Clinical Care
49	431	Preventive Care and Screening Unhealthy Alcohol Use: Screening & Brief Counseling	Percentage of patients aged 18 years and older who were screened at least once within the last 24 months for unhealthy alcohol use using a systematic screening method AND who received brief counseling if identified as an unhealthy alcohol user	Yes	Process	Community/ Population Health
50	438	Statin Therapy for the Prevention and Treatment of Cardiovascular Disease	Percentage of the following patients – all considered at high risk of cardiovascular events – who were on statin therapy during the performance period: <ul style="list-style-type: none"> <li>Adults aged <math>\geq 21</math> years who were previously diagnosed with or currently have an active diagnosis of clinical atherosclerotic cardiovascular disease (ASCVD); OR</li> <li>Adults <math>\geq 21</math> years who ever had a fasting or direct low-density lipoprotein cholesterol (LDL-C) level <math>\geq 190</math> mg/dl or were previously diagnosed with or currently have an active diagnosis of familial or pure hypercholesterolemia; OR</li> <li>Adults aged 40-75 years with a diagnosis of diabetes with a fasting or direct LDL-C level of 70-189 mg/dl</li> </ul>	Yes	Process	Effective Clinical Care
51	eCQM* CMS347v3			Yes		



**QPP Measures**

#	QPP #	Measure Title	Definition	Historical Benchmark?	Type	Domain
52	<b>462</b> <i>eCQM*</i> <i>CMS645v3</i> <i>version only</i>	Bone density evaluation for patient with Prostate Cancer and receiving Androgen Deprivation Therapy	Patients determined as having prostate cancer who are currently starting or undergoing androgen deprivation therapy (ADT), for an anticipated period of 12 months or greater and who receive an initial bone density evaluation. The bone density evaluation must be prior to the start of ADT or within 3 months of the start of ADT	Yes	Process	Effective Clinical Care
53	<b>476</b> <i>eCMQ*</i> <i>CMS771v1</i> <i>version only</i>	Urinary Symptom Score Change 6-12 Months After Diagnosis of Benign Prostatic Hyperplasia	Percentage of patients with an office visit within the measure period and with a new diagnosis of clinically significant Benign Prostatic Hyperplasia who have an International Prostate Symptom Score (IPSS) or American Urological Association (AUA) Symptom Index (SI) documented at the time of diagnosis and again 6-12 months later with an improvement of 3 points	Yes	Outcome	Person and Caregiver- Centered Experience and Outcomes