

PTNS References

The following highlight clinical use of PTNS as a treatment for Overactive Bladder (OAB) and associated symptoms of urinary frequency, urinary urgency and urge incontinence.

2012

Burton, C., Sajja, A., Latthe, P.M. (2012). Effectiveness of percutaneous posterior tibial nerve stimulation for overactive bladder: a systematic review and meta-analysis. *Neurourology and Urodynamics*, epub May 15.

Monga, A., Tracey, M., Subbaroyan, J. (2012). A systematic review of clinical studies of electrical stimulation for treatment of lower urinary tract dysfunction. *Int Urogynecol J*, epub March 20.

Peters, K., Carrico, D., MacDiarmid, S., Wooldridge, L., Khan, A., McCoy, C. et al. (2012). Sustained therapeutic effects of percutaneous tibial nerve stimulation: 24-month results of the STEP study. *Neurourology and Urodynamics*, epub June 5.

Staskin, D., Peters, K., MacDiarmid, S., Shore, N., de Groat, W. (2012). Percutaneous tibial nerve stimulation: a clinically and cost effective addition to the overactive bladder algorithm of care. *Curr Urol Rep*, epub August 15.

2011

Leong, F., McLennan, M.T., Barr, S.A., & Steele, A.C. (2011). Posterior tibial nerve stimulation in patients who have failed anticholinergic therapy: efficacy and time to response. *Female Pelvic Med Reconstr Surg*, 17(2), 74-75.

Ruan, J., Dahabreh, I., Ip, S., Ferzandi, T., & Wong, J. (2011). Percutaneous tibial nerve stimulation in the management of overactive bladder: a systematic review and meta-analysis. *Female Pelvic Med Reconstr Surg*, 17(5), supplement 2.

2010

Finazzi-Agrò, E., Petta, F., Sciobica, F., Pasqualetti, P., Musco, S., & Bove, P. (2010). Percutaneous tibial nerve stimulation effects on detrusor overactivity incontinence are not due to a placebo effect: a randomized, double-blind, placebo controlled trial. *J Urol*, 184, 2001-2006.

MacDiarmid, S.A., Peters, K.M., Shobeiri, S.A., Wooldridge, L.S., Rovner, E.S., Leong, F.C., et al. (2010). Long-term durability of percutaneous tibial nerve stimulation for the treatment of overactive bladder. *J Urol*, 183, 234-240.

Peters, K.M., Carrico, D.J., Perez-Marrero, R.A., Khan, A.U., Wooldridge, L.S., Davis, G.L., & MacDiarmid, S.A. (2010). Randomized trial of percutaneous tibial nerve stimulation versus sham efficacy in the treatment of overactive bladder syndrome: Results from the SUMIT Trial. *J Urol*, 183, 1438-1443.

Ridout, A.E., & Yoong, W. (2010). Tibial nerve stimulation for overactive bladder syndrome unresponsive to medical therapy. *Journal of Obstetrics and Gynaecology*, 30(2), 111-114.

2009

MacDiarmid, S.A., & Staskin, D.R. (2009). Percutaneous tibial nerve stimulation (PTNS): A literature-based assessment. *Current Bladder Dysfunction Reports*, 4, 29-33.

Peters, K.M., MacDiarmid, S.A., Wooldridge, L.S., Leong, F.C., Shobeiri, S.A., Rovner, E.S., et al. (2009). Randomized trial of percutaneous tibial nerve stimulation versus extended-release tolterodine: Results from the overactive bladder innovative therapy trial. *J Urol*, 182, 1055-1061.

Wooldridge, L.S. (2009). Percutaneous tibial nerve stimulation for the treatment of urinary frequency, urinary urgency, and urge incontinence: Results from a community-based clinic. *Urologic Nursing*, 29(3), 177-185.

Zinkgraf, K., O'Leary Quinn, A., Ketterhagen, D., Kreuziger, B., & Stevenson, K. (2009). Percutaneous Tibial Nerve Stimulation for Treatment of Overactive Bladder and Urinary Retention in an Elderly Population. *Urologic Nursing*, 29(1), 30-34.

2007

Farahat, Y.A. (2007). Prospective randomized study for long term efficacy of percutaneous tibial nerve stimulation (PTNS) in patients with over active bladder: Maintenance protocol. Abstract, 4th Pan Arab Continence Society Conference, Qatar.

Finazzi-Agrò, E., Capellano, F., Miano, R., Catanzaro, M., Petta, F., Sciobica, F., et al. (2007). Who are the best candidates to be treated with percutaneous tibial nerve stimulation (PTNS)? A retrospective analysis of responder rate in different patients' populations. *European Urology Supplement*, 6(2), 141.

Preyer, O., Gabriel, B., Mailath-Pokony, M., Doerfler, D., Laml, T., Umek, W., et al. (2007). Peripheral tibial neurostimulation (PTNS) versus Tolterodine in the treatment of women with urge urinary incontinence and urge symptoms. *Intern Urogynecol J, Suppl 1*.

Van Balken, M. (2007). Percutaneous tibial nerve stimulation: The Urgent PC device. *Expert Review of Medical Devices*, 4(5), 693-698.

2006

Finazzi-Agrò, E., Petta, F., Pachatz, C., Miano, R., Sciobica, F., & Marfia, G.A. (2006). Long latency somatosensory evoked potentials: Modifications after percutaneous tibial nerve stimulation. *European Urology*, 5(2), 259.

van Balken, M.R., Vergunst, H., & Bemelmans, B.L.H. (2006). Prognostic factors for successful percutaneous tibial nerve stimulation. *European Urology*, 49, 360-365.

van der Pal, F., van Balken, M.R., Heesakkers, J.P.F.A., DeBruyne, F.M.J., & Bemelmans, B.L.H. (2006). Percutaneous tibial nerve stimulation in the treatment of refractory overactive bladder syndrome: Is maintenance treatment necessary? *BJU International*, 97, 547-550.

van der Pal, F., van Balken, M.R., Heesakkers, J.P.F.A., DeBruyne, F.M.J., Kiemeny, L.A.L.M., & Bemelmans, B.L.H. (2006). Correlation between quality of life and voiding variables in patients treated with percutaneous tibial nerve stimulation. *BJU International*, *97*, 113-116.

2005

Cooperberg, M.R., & Stoller, M.L. (2005). Percutaneous neuromodulation. *Urology Clinics of North America*, *32*, 71-78.

Finazzi-Agró, E., Campagna, A., Sciobica, F., Petta, F., Germani, S., Zuccalà, A., et al. (2005). Posterior tibial nerve stimulation (PTNS): Is the once-a-week protocol the best option? *Minerva Urol Nefrol*, *57*(2), 119-123.

Karademir, K., Baykal, K., Sen, B., Senkul, T., Iseri, C., & Erden, D. (2005). A peripheral neuromodulation technique for curing detrusor overactivity: Stoller afferent neurostimulation. *Scandinavian Journal of Urology and Nephrology*, *39*, 230-233.

Nuhoglu, B., Fidan, V., Ayyildiz, A., Karaguzel, E., Huri, E., Ersoy, E., et al. (2005). Stoller Afferent Nerve Stimulation (SANS) in women with therapy resistant overactive bladder: A 1-year follow-up. *European Urology*, *3*, 123.

van der Pal, F., van Balken, M.R., Heesakkers, J.P.F.A., DeBruyne, F.M.J., & Bemelmans, B.L.H. (2005). Maintenance percutaneous tibial nerve stimulation (PTNS) treatment in successfully treated patients with refractory overactive bladder syndrome a necessity. *European Urology*, *3*, 144.

Vincente, E., Hannaoui, N., Gonzalez, J.L., Pena, J.A., Prats, J., Garcia, D., et al. (2005). Posterior tibial neuromodulation (PTNM) in the treatment of overactive bladder (OAB). *International Journal Proctological Perineal Disease*, *IX*(1), 31, A29.

2004

Ruiz, C.B., Outeiriño, P.X.M., Martínez, P.C., Dueñas, E.L., & López, A.L. (2004). Peripheral afferent nerve stimulation for treatment of lower urinary tract irritative symptoms. *European Urology*, *45*, 65-69.

van Balken, M.R., Vergunst, H., & Bemelmans, B.L.H. (2004). The use of electrical devices for the treatment of bladder dysfunction: A review of methods. *Journal of Urology*, *172*, 846-851.

Vandoninck, V., van Balken, M.R., Finazzi-Agró, E., Heesakkers, J.P.F.A., DeBruyne, F.M.J., Kiemeny, L.A.L.M., et al. (2004). Posterior tibial nerve stimulation in the treatment of voiding dysfunction: Urodynamic data. *Neurourology and Urodynamics*, *23*, 246-251.

2003

Vandoninck, V., van Balken, M.R., Finazzi-Agró, E., Petta, F., Caltagirone, C., Heesakkers, J.P.F.A., et al. (2003). Posterior tibial nerve stimulation in the treatment of urge incontinence. *Neurourology and Urodynamics*, *22*, 17-23.

Vandoninck, V., van Balken, M.R., Finazzi-Agró, E., Petta, F., Micali, F., Heesakkers, J.P.F.A., et al. (2003). Percutaneous tibial nerve stimulation in the treatment of overactive bladder. Urodynamic data. *Neurourology and Urodynamics*, *22*, 227-232.

Vandoninck, V., van Balken, M.R., Finazzi-Agró, E., Petta, F., Micali, F., Heesakkers, J.P.F.A., et al. (2003). Posterior tibial nerve stimulation in the treatment of idiopathic nonobstructive voiding dysfunction. *Urology*, *61*(3), 567-572.

2002

Mazo, E.B., & Krivoborodov, G.G. (2002). Temporary sacral and tibial neuromodulation in treating patients with overactive urinary bladder. *Zurnal Voprosy Neirochirurgii Imeni NN Burdenko*, *1*, 17-21.

2001

Govier, F.E., Litwiller, S., Nitti, V., Kreder, K.J. Jr., & Rosenblatt, P. (2001). Percutaneous afferent neuromodulation for the refractory overactive bladder: Results of a multi-center study. *Journal of Urology*, *165*, 1193-1198.

van Balken, M.R., Vandoninck, V., Gisolf, K.W.H., Vergunst, H., Kiemeny, L.A.L.M., DeBruyne, F.M.J., et al. (2001). Posterior tibial nerve stimulation as neuromodulative treatment of lower urinary tract dysfunction. *Journal of Urology*, *166*, 914-918.

2000

Finazzi-Agró, E., Petta, F., Giannantoni, A., D'Amico, A., Germani, S., Bitelli, M., et al. (2000). Use of Stoller Afferent Nerve Stimulation (SANS) in the treatment of detrusor hyperactivity: Preliminary data. *Urologica*, *10*, 155-156.

Fiori, M., Mercuriali, M., Fabbri, F., Gunelli, R., Saltutti, C., Trinci, E., et al. (2000). SANS (Stoller Afferent Nerve Stimulation) a new option in the treatment of overactive bladder. *Urologica*, *10*, 157-158.

Klingler, H.C., Pycha, A., Schmidbauer, J., & Marberger, M. (2000). Use of peripheral neuromodulation of the S3 region for treatment of detrusor overactivity: A urodynamic-based study. *Urology*, *56*, 766-771.

1999

Craggs, M., & McFarlane, J. (1999). Physiological Society Symposium: The physiology and pathophysiology of the urinary tract: Neuromodulation of the lower urinary tract. *Experimental Physiology*, *84*, 149-160.

Govier, F.E., Litwiller, S.E., & Mitchell, I. (1999). Percutaneous peripheral nerve stimulation for urgency/frequency syndrome. *European Urology*, *35*(52), 16.

Stoller, M.L. (1999). Nerve stimulation for pelvic floor dysfunction. *European Urology*, *35*(suppl 2), 16.